

Patient Instructions for Heart Catheterization and Balloon Angioplasty Instructions

Patien	t Name:						
		Your prod	cedure is	scheduled v	<u>with:</u>		
Dr. Lieber	Dr. Uddin	Dr. Raza	Dr. Wen	Dr. Salazar	Dr. Bellera	Dr. Almahm	oud
Date:							
Arrival Time:		☐ AM [] РМ	Procedure Ti	me:	Д АМ	🔲 РМ
We cannot predict occur, or that the coccupy your time in	ase ahead	of you may	-		•	-	-
You will need to pro and EKG if necess registering yourself	ary. Please I	oring your i		•			•
Your pre-registrat	ion is sched	duled on: _					
Arrive at:		и 🔲 РМ					
Humble Surgic	al Hospital [A hospital	representa	tive will call ye	ou to schedu	le]	
Kingwood Med	lical Center	[Entrance (C, Suite 20	0 Phone: (28	31) 348-8344]		
Memorial Herm	nan Northea	st Hospital	[Southwes	st Tower Pho	ne: (281) 540)-7936]	
Houston North	west Hospit	al [Medica	l Mall II Ph	none: (281) 44	0-2600]		
]	Day Of Yo	our Proced	dure Instruc	<u>tions</u>		
 Nothing to eat of 	or drink the	night befor	e your prod	cedure			

- Take your routine medication with a glass of water the morning of your procedure
- You will need someone to drive you home
- Patients on Insulin: You will need to check your blood sugar in the morning. If it is under 200, hold your insulin. If it's between 200-400, take half of your normal morning dose. If it's over 400 take your whole morning dose of insulin. Please bring your glucometer and insulin with you to the hospital, so you can manage your own blood sugar levels and dosages throughout the day.
- ◆ Patients on Glucophage/Metformin: STOP Glucophage/Metformin 1 day prior to your procedure and resume 48 hours after your procedure.
- Patients on Aspirin, Plavix, Effient or Brilinta: Continue these medications. If anyone requests you discontinue these medications, please contact our office.

tcahouston.com **LAST UPDATED 1/14/2021**



Information on Nuclear Stress Test 1/2

Day Of Your Procedure Instructions

- ◆ Patients on Coumadin, Eliquis or Pradaxa: STOP 5 days prior to your procedure and resume 1 day after your procedure.
- ◆ Patients on Xarelto: STOP medication 24 hours prior to your procedure and resume one day after your procedure.
- ◆ <u>DO YOU HAVE ANY ALLERGIES TO SHELLFISH OR IODINE?</u>
 If you answered yes to this question, then you will need to be pre-medicated 2 days before your procedure for prophylactic reasons.

ABOUT THE PROCEDURE

The nurse will need to start an IV. They may need to shave your groin area where the doctor will be inserting the catheter. Before they can insert the catheter, they will inject local anesthesia into the groin site. This is the only pain you will feel. You will get medication to sedate you for the procedure (Valium, Benadryl and/or Versed). You will be awake during the procedure will be able to talk to the physician while the procedure is in progress. When they inject the dye, you may feel a warm flushing sensation, but you should not feel any pain. After the doctor removes the catheter they will then place a "plug" in the artery.

ABOUT THE PROCEDURE

Once the "plug" is in place you will need to be on bed rest for the following 2-3 hours. If you only have a heart catheterization, you will be able to go home in the evening. If you have a balloon angioplasty or stent placement, you will have an overnight stay in the hospital and will be released the next day. Some bruising and/or a knot to your groin area is normal. Call if you encounter increasing pain, fever, redness or discharge at the incision site. For the first 2 days following the procedure, no lifting, bending or squatting. You should be able to return to work following your procedure within 2-3 days. You may shower and remove your dressing the next day following your procedure. Keep incision site covered with a Band-Aid after dressing removal for 3-5 days. Do not wear anything binding over and/or around the incision.

Your hospital follow-up is scheduled with:

Date:		Appo	ointment Tim	ne:	AM 🔲 PM			
Dr. I	Lieber	Dr. Uddin	Dr. Raza	Dr. Wen	Dr. Salazar	Dr. Bellera	Dr. Almahmo	oud

If you have any questions, please call our office at (281)-358-1950